

SYSTEMATIC REVIEW OF CLINICAL DECISION SUPPORT INTERVENTIONS  
WITH POTENTIAL FOR INPATIENT COST REDUCTION

by

Christopher Fillmore

A thesis submitted to the faculty of  
The University of Utah  
in partial fulfillment of the requirements for the degree of

Master of Science

Department of Biomedical Informatics

The University of Utah

August 2013

Copyright © Christopher Fillmore 2013

All Rights Reserved

# The University of Utah Graduate School

## STATEMENT OF THESIS APPROVAL

The thesis of **Christopher Fillmore**

has been approved by the following supervisory committee members:

|                         |          |                     |
|-------------------------|----------|---------------------|
| <b>Kensaku Kawamoto</b> | , Chair  | <b>7 June 2013</b>  |
| <hr/>                   |          | <hr/> Date Approved |
| <b>Bruce Bray</b>       | , Member | <b>7 June 2013</b>  |
| <hr/>                   |          | <hr/> Date Approved |
| <b>Charlene Weir</b>    | , Member | <b>7 June 2013</b>  |
| <hr/>                   |          | <hr/> Date Approved |

and by **Julio Facelli**, Chair of  
the Department of **Biomedical Informatics**

and by Donna M. White, Interim Dean of The Graduate School.

## **ABSTRACT**

The purpose of this thesis was to systematically review trials of clinical decision support interventions with the potential to reduce inpatient costs, so as to identify promising interventions for more widespread implementation and to inform future research in this area. MEDLINE was searched up to September 2012, and relevant studies were identified using titles and abstracts. Full text articles were reviewed to make a final determination on inclusion. Relevant characteristics of the studies were extracted and summarized. Following a screening of 6,978 articles, 60 manuscripts were included. The majority of manuscripts were published during or after 2007. 63.3% of studies were pre-post comparisons, and 13.3% were randomized controlled trials. 56.7% of the studies were focused on pharmacotherapy. 71.7% of the studies resulted in statistically and clinically significant improvements in an explicit financial measure or a proxy financial measure. Only 15% of the studies directly measured the financial impact of an intervention, whereas the financial impact was inferred in the remainder of studies. Data on cost-effectiveness were available for only one study. Given these results, it is apparent that further research is needed on the cost impact and cost effectiveness of CDS in the inpatient setting.

## TABLE OF CONTENTS

|   |     |
|---|-----|
| ABSTRACT .....                          | iii |
| ACKNOWLEDGEMENTS .....                  | v   |
| BACKGROUND .....                        | 1   |
| METHODS .....                           | 3   |
| Data Source .....                       | 3   |
| Inclusion and Exclusion Criteria .....  | 3   |
| Study Selection .....                   | 3   |
| Data Extraction .....                   | 4   |
| Data Analysis .....                     | 4   |
| RESULTS AND DISCUSSION.....             | 5   |
| Study Timing.....                       | 5   |
| Study Designs .....                     | 5   |
| Study Quality .....                     | 6   |
| Clinical Focus .....                    | 8   |
| Cost Effectiveness .....                | 8   |
| Direct Measurement of Cost .....        | 9   |
| Use of Proxy Measures .....             | 10  |
| Improvement in Cost/Proxy Measures..... | 12  |
| CONCLUSIONS.....                        | 32  |
| APPENDIX .....                          | 33  |
| REFERENCES .....                        | 34  |

## **ACKNOWLEDGEMENTS**

This work was supported in part by National Library of Medicine training grant T15-LM07124.

## **BACKGROUND**

Healthcare costs are increasing rapidly and at an unsustainable rate in many countries. In the United States, inpatient care is the single largest contributor to national health expenditures, accounting for 31.5% of \$2.7 trillion dollars of health expenditures in 2011 [1]. As such, inpatient care is a significant driver of increased health spending. In 2011, the annual spending on hospital care in the U.S. grew 4.3% as compared to 3.9% growth in overall health expenditures [1]. Contributing to the importance of addressing inpatient costs is the fact that reducing these costs has the potential to financially benefit inpatient healthcare organizations regardless of reimbursement models. Traditional episode-of-care payment systems (for example, Medicare's inpatient prospective payment system), bundled payments systems, and comprehensive payment systems (embodied in accountable care organizations) are all examples of reimbursement models under which healthcare organizations stand to benefit from reducing inpatient costs [2].

Clinical decision support (CDS) represents a promising approach to both improving outcomes and decreasing costs [3]. Several past systematic reviews have examined outcomes related to clinical decision support systems in the inpatient setting, but few have focused on the impact of CDS on inpatient costs specifically [4-9]. One review published in 2006 evaluated cost as an outcome [10]. However, this review was focused on health information technology (IT) in general rather than CDS specifically. Moreover, it only included studies published through January 2004. A second, more

recent review on CDS included cost outcomes but was limited to studies with a randomized trial design [11]. The design and timing of these two reviews potentially excluded relevant CDS intervention trials. In particular, nonrandomized research designs are commonly used to evaluate CDS interventions.

Given the importance of limiting the growth of inpatient costs and the potential benefit of CDS, we sought to (i) inclusively identify promising interventions that could serve as models for more widespread implementation and to (ii) identify gaps in the literature warranting further research. As such, we systematically reviewed both randomized and nonrandomized trials of CDS systems with the potential to reduce inpatient or emergency department (ED) costs.



## **METHODS**

### **Data source**

Using a search strategy adapted from a previous systematic review [7], we searched MEDLINE through September 24, 2012. The latest search was performed on that date. We used a combination of the following search terms: *decision support systems, clinical; decision-making, computer-assisted; reminder systems; guideline adherence; and medical informatics*. Details of the search strategy are available in the Appendix. Search results were limited to human subjects and the English language.

### **Inclusion and Exclusion Criteria**

We defined a CDS system as a system designed to directly aid in clinical decision making, in which characteristics of individual patients are matched to a knowledge base for the purpose of presenting patient-specific assessments or recommendations to clinicians [4]. Inclusion criteria were as follows: peer-reviewed primary manuscript; clinical trial of a CDS system in an inpatient or ED setting; and use of either cost or a proxy measure for cost (e.g., length of stay) as an outcome metric. Exclusion criteria were as follows: study not in English; or use of CDS in the control group.

### **Study Selection**

Titles and abstracts from retrieved references were reviewed to determine potential inclusion eligibility. The full texts of studies that appeared to be potentially

eligible were then reviewed. Final inclusion determinations were made using the full texts. In cases where a study's inclusion status was unclear upon review by the primary reviewer, the authors jointly reviewed the study and made a consensus decision.

### **Data Extraction**

For each article that met inclusion criteria, data were extracted on setting, trial design, intervention, and trial results. Abstracted trial results included outcomes with potential cost-saving implications, whether costs were directly measured, whether there was a statistically and clinically significant improvement in cost or in a proxy measure, and whether the study could be considered a cost-effectiveness study. Specifically, measures with potential cost-saving implications consisted of direct cost measures or other proxy measures with cost ramifications. Proxy cost measures included length of stay, adverse events, and process measures correlated with adverse events. Clinical significance of results was determined by author consensus. To be considered a cost effectiveness study, the study must have accounted at least for the personnel costs included in developing and deploying the intervention. For commercial CDS systems, at least the cost of licensing the software must also have been considered.

### **Data Analysis**

Extracted data were analyzed and presented in table form and narrative summary. Additionally, significant themes, trends, and patterns were noted and discussed.

## **RESULTS AND DISCUSSION**

The literature search returned a total of 6,978 unique references. 6,855 references were excluded after screening of titles and abstracts. We reviewed 123 full-text articles, of which 60 [12-71] met criteria for inclusion in the review (see Figure 1). Characteristics of these studies are summarized in Table 1.

### **Study Timing**

A majority (55%) of studies were published during or after 2007 [39-71]. The earliest included study was published in 1989 [12]. This high concentration of studies published between 2007 and 2012 represents large recent growth in the evaluation of inpatient CDS systems and is consistent with increasing adoption of health IT generally.

### **Study Designs**

Almost all (86.7%) [12-19, 21-23, 25-33, 35, 36, 38, 39, 41-43, 45-59, 61-63, 65-71] of the studies were quasi-experimental trials (i.e., studies that aim to evaluate interventions but that do not use randomization [72]). Overall, the most common design used in the studies was a pre-post comparison, wherein researchers used historical controls prior to the implementation of an intervention. Of the 60 studies, 40 (66.7%) were some form of pre-post comparison [12, 14-16, 18, 25-27, 29-33, 36, 38, 39, 41, 42,

46-59, 61-63, 66-68, 70, 71]. Only eight (13.3%) of the studies were randomized controlled trials [20, 24, 34, 37, 40, 44, 60, 64].

The frequent use of quasi-experimental designs in medical informatics evaluations has been noted previously [72, 73], and the results of this review are consistent. Given the overwhelming prevalence of quasi-experimental designs, reviews of CDS systems that only include RCTs are bound to exclude a large portion of the published literature. With respect to this systematic review, a deliberate decision was made to include quasi-experimental studies, as one of the primary goals of this study was to develop a comprehensive catalog of CDS interventions that have the potential for reducing inpatient costs. At the same time, the inclusion of quasi-experimental study designs may have resulted in the inclusion of studies that are more subject to bias than RCTs.

### **Study Quality**

Given that the stated goals of this review are exploratory in nature, we opted not to perform a meta-analysis of the extracted data. Moreover, because we chose to be inclusive in study selection, the studies in the review were heterogeneous in terms of interventions, outcomes, and study design. Additionally, the actual number of studies that included explicit economic data was very small. These factors limit the extent to which a meaningful synthesis of data might be obtained from meta-analysis techniques.

However, under different circumstances if a meta-analysis were performed, it would be customary to evaluate the quality of the included studies to provide an indication as to the validity of aggregate conclusions. In a review where the study designs are primarily quasi-experimental (such as this review), certain factors make the

evaluation of quality difficult. One factor is the number of different types of trial designs that fall under the quasi-experimental category. Within the domain of medical informatics, one review identified eleven different quasi-experimental trial designs [72]. These designs fell under the following broad categories: (a) quasi-experimental designs without control groups, (b) quasi-experimental designs that use a control group but no pretest, (c) quasi-experimental designs that use control groups and pretests, and (d) interrupted time-series designs [72]. Presumably, different trial designs increase the difficulty of the evaluation process. Notably, the Cochrane Collaboration recommends that systematic reviews that include nonrandomized studies employ a dedicated methodologist to assist in the evaluation of bias within studies due to the difficulty of addressing multiple study designs [74].

A second factor potentially hindering this evaluation is the lack of widely accepted quality assessment tools for evaluation of nonrandomized intervention studies. One systematic review on this topic identified 194 tools that have been used to assess the quality of nonrandomized intervention studies [75]. However, the authors of this review concluded that only 6 of these tools were appropriate for use in systematic reviews and that most of these tools would benefit from further modification. If we were to perform a quality analysis in the current review, an appropriate tool might be the Quality Assessment Tool for Quantitative Studies [76]. This tool was initially developed for evaluation of studies in the public health domain where nonrandomized study designs are commonly used. The tool evaluates 6 primary components: selection bias, design, confounders, blinding, data collection methods, and withdrawals/dropouts [76]. The tool

is reportedly easy to use, is not time consuming, and has comprehensive documentation available to assist with use [75].

### **Clinical Focus**

The most common clinical focus targeted by CDS systems in the review was pharmacotherapy, with 56.7% of studies focused on this area [12, 13, 15, 16, 19, 23, 25-27, 29-33, 35-38, 41, 42, 46, 47, 50, 51, 54, 55, 59, 62-66, 69, 71]. The second most common area of clinical focus was venous thromboembolism prophylaxis, which accounted for 11.7% of studies in the review [20, 21, 24, 34, 43, 44, 53]. Examples of other clinical areas addressed included blood transfusion management [39, 40, 57], sepsis management [56, 61], and heparin-induced thrombocytopenia diagnosis [49, 58].

The significant focus on pharmacotherapy within the included studies may reflect the importance of drug selection within computerized provider order entry (CPOE) systems, which are foundational to CDS in many inpatient settings. 51.7% [15, 16, 19, 21, 23, 24, 26, 27, 29-34, 36, 39-41, 47, 49, 50, 53, 54, 57, 59, 62, 65, 68-71] of the studies overall involved CDS in the context of CPOE, and 61.8% [15, 16, 19, 23, 26, 27, 29-33, 36, 41, 47, 50, 54, 59, 62, 65, 69, 71] of the pharmacotherapy studies involved CDS in this context.

### **Cost-Effectiveness**

Only one of the 60 studies (1.7%) was considered to be a cost-effectiveness study [52]. This study evaluated the use of a well-known diagnostic decision support system, DXplain, with residents in a teaching hospital. The authors reported that access to DXplain had been provided at no charge for the purposes of the study, but that an annual

license would have cost their organization \$4,000-\$6,000 per year [52]. It is telling that the only study to address cost effectiveness in this review concerned a simple license to a stand-alone, diagnostic CDS system. The majority of the studies in this review dealt with more comprehensive, integrated systems either purchased through vendors or developed locally. Under those circumstances, providing information about cost of development, implementation, or licensing fees is presumably more difficult. However, the near complete lack of this type of information is concerning given the need for such cost-effectiveness information by public policy developers and decision makers within healthcare organizations.

### **Direct Measurement of Cost**

Nine (15%) studies in the review directly measured costs [16, 23, 26, 27, 29, 37, 39, 42, 52]. Seven of these studies focused on pharmacotherapy [16, 23, 26, 27, 29, 37, 42], while the other two addressed management of blood transfusion [39] and general medical diagnosis [52]. These studies included one RCT [37], with the remainder of studies having a quasi-experimental design. Of these studies, four reported a statistically and clinically significant improvement in a cost measure [16, 37, 39, 52]. Except in one case as outlined, the cost involved in implementing these interventions was not studied. Because investments in CDS are like any other business investment, having only one side of the financial equation (cost impact) is insufficient for making public policy and business decisions.

### **Use of Proxy Cost Measures**

85% of the studies in the review solely reported proxy measures as indicators of impact on cost [12-15, 17-22, 24, 25, 28, 30-36, 38, 40, 41, 43-51, 53-71]. The most commonly used type of proxy measure in this group was process measures that were associated with adverse events. Of the studies that solely used proxy measures, 60.8% reported this type of measure [13, 15, 17, 19, 21, 24, 25, 30, 32, 33, 35, 36, 41, 45-50, 54-56, 58-60, 62-64, 66, 68, 71]. Examples of other proxy measures reported by these studies included rates of adverse events (reported by 21.6% of studies [12, 15, 34, 43, 51, 53, 59, 65, 68, 69, 71]), length of stay (reported by 9.8% of studies [20, 44, 60, 61, 65]), and patient charges (reported by 5.9% of studies [14, 22, 28]).

As noted, a strikingly small percentage of the studies directly measured an intervention's impact on cost. Therefore, in the majority of cases, we were left to infer a possible cost savings from nonfinancial proxy measures. Doing so has some inherent limitations. For example, four studies reported patient charges as an outcome [14, 22, 28, 52]. This is not a direct measure of cost, and it can be unclear as to how charges actually relate to cost [77]. We assumed that an institution's costs were at least proportional to what it charged a patient. However, given that we did not know the actual relationship between costs and charges at any given institution, this assumption suffered from an element of uncertainty.

Other limitations of using proxy measures for cost were apparent in our data. Two studies reported no differences in actual measured costs but reported decreases in length of stay [23, 27]. For this review, we considered length of stay a reasonable proxy measure for cost. However, in these two instances, shorter lengths of stay did not coincide with



actual decreased costs. The reverse of this situation was present in two studies, where directly measured costs decreased, but no difference in length of stay was detected [37, 52]. It is notable that of the nine studies in the review that directly measured costs, four demonstrated discrepancies between explicit cost measures and available proxy cost measures.

Another limitation of using proxy measures is related to adverse events. We considered measures of adverse events an appropriate proxy measure for cost given the potential for these events to necessitate the utilization of additional resources. We went a step further and included process measures correlated with adverse events as proxy measures as well. For example, one study in the review reported the rate of compliance with venous thromboembolism prophylaxis guidelines (a process measure correlated with an adverse event) [21]. Alternatively, another study reported the actual incidence of venous thromboembolism (a measure of an adverse event) [34]. For this review, we made the assumption that an improvement in a process measure associated with an adverse event would be associated with an improvement in the incidence of that adverse event. Decreased incidence of an adverse event, in turn, would be associated with cost savings. However, in one study, process measures correlated with an adverse event were significantly improved, but there was no improvement in the incidence of the actual adverse event [59]. More perplexingly, another study reported improvements in a process measure, no improvement in the correlated adverse event, and a significant improvement in length of stay [65].

### **Improvement in Cost/Proxy Measures**

43 (71.7%) of the studies reported a statistically and clinically significant improvement in a cost or proxy measure [12, 13, 15, 16, 18, 19, 21, 24, 28, 30-40, 43, 45-47, 51, 52, 54-60, 62-71]. However, when considered in the context of the lack of direct cost measurements, the limitations of proxy cost measures, and the prevalence of quasi-experimental designs, it is difficult to know what level of confidence to place in that finding. On the face of it, CDS does appear to be a promising intervention for reducing inpatient costs. However, further research is clearly needed in order to more concretely characterize the benefits that have been achieved and that might be achieved in the future.




Figure 1, Study Selection Flow Chart

Table 1, Study Characteristics

| <i>Citation</i>       | <i>Setting</i>                | <i>Scope</i>                         | <i>Trial design</i>     | <i>Intervention (control = usual care unless otherwise specified)</i>   | <i>Measures with potential cost saving implications</i>                                     | <i>Results (intervention vs. control)</i>          | <i>Direct measurement of cost</i> | <i>Integrated with primary clinical information system</i> | <i>Statistically &amp; clinically significant improvement in cost or proxy metric</i> | <i>Cost effectiveness study</i> |
|-----------------------|-------------------------------|--------------------------------------|-------------------------|---|---|--|-----------------------------------|--|---|---------------------------------|
| Larsen RA (1989) [12] | Teaching hospital             | 1 hospital; 6831 patients            | Pre-post comparison     | Computer generated paper reminders of need for perioperative antibiotics  | Surgical wound infection rates  | 0.9% vs. 1.8% (P < 0.03)                           | No                                | Yes  | Yes   | No                              |
| Rind DM (1991) [13]   | Teaching hospital             | 1 hospital; 10076 patients           | Interrupted time series | CDS designed to alert physicians to rising creatinine levels in patients taking nephrotoxic drugs   | Mean time from rise in creatinine to discontinuation or dose adjustment of nephrotoxic drug | 72.6 vs. 93.7 hours (P < 0.001)                    | No                                | Yes  | Yes   | No                              |
| Day F (1995) [14]     | Academic emergency department | 1 emergency department; 465 patients | Pre-post comparison     | Guideline-driven electronic charting system that provided recommendations for evaluation and treatment of patients with acute low back pain | Mean charges  | \$257 vs. \$239 (NS)                               | No                                | No   | No  | No                              |
| Bates DW (1998) [15]  | Academic hospital             | 1 hospital; 6711 admissions          | Pre-post comparison     | CPOE with CDS   | Non-intercepted serious medication error rate   | 4.86 vs. 10.7 events/1000 patient days (P = 0.01)  | No                                | Yes  | Yes   | No                              |
|                       |                               |                                      |                         |   | Preventable ADE rate  | 3.88 vs. 4.69 events/1000 patient days (NS)        |                                   |  |   |                                 |
|                       |                               |                                      |                         |   | Non-intercepted potential ADE rate  | 0.98 vs. 5.99 events/1000 patient days (P = 0.002) |                                   |  |   |                                 |

Table 1 Continued

| <i>Citation</i>        | <i>Setting</i>                | <i>Scope</i>              | <i>Trial design</i>                    | <i>Intervention (control = usual care unless otherwise specified)</i>   | <i>Measures with potential cost saving implications</i>                          | <i>Results (intervention vs. control)</i> | <i>Direct measurement of cost</i> | <i>Integrated with primary clinical information system</i> | <i>Statistically &amp; clinically significant improvement in cost or proxy metric</i> | <i>Cost effectiveness study</i> |
|------------------------|-------------------------------|---------------------------|--|---|--|---|-----------------------------------|--|---|---------------------------------|
| Evans RS (1998) [16]   | Teaching hospital             | 1 hospital; 1681 patients | Pre-post comparison                    | Comprehensive antimicrobial management system in an ICU setting   | Mean total cost of hospitalization when computer regimen was followed            | \$26315 vs. \$35283 (P < 0.001)           | Yes                               | Yes  | Yes   | No                              |
|                        |                               |                           |  |   | Mean cost of anti-infective agents when computer regimen was followed            | \$102 vs. \$340 (P < 0.001)               |                                   |  |   |                                 |
|                        |                               |                           |  |   | Total hospital length of stay when computer regimen was followed                 | 10.0 vs. 12.9 days (P < 0.001)            |                                   |  |   |                                 |
| Knirsch CA (1998) [17] | Academic hospital             | 1 hospital; 43 patients   | Prospective evaluation without control | Automated computer protocol for identification of patients requiring tuberculosis isolation combined with a clinical protocol | Isolation of culture positive tuberculosis patients within 24 hours of admission | 79% vs. 70% (NS)                          | No                                | No   | No  | No                              |
| Aase O (1999) [18]     | Academic emergency department | 1 ED; 493 patients        | Pre-post comparison                    | CDS using Bayes theorem to assist in diagnosis and triage of patients with acute chest pain                                   | Percent of patients unnecessarily admitted to coronary care unit                 | 19% vs. 35% (P < 0.05) <sup>#</sup>       | No                                | No   | Yes <sup>#</sup>  | No                              |

Table 1 Continued

| <i>Citation</i>         | <i>Setting</i>                   | <i>Scope</i>                | <i>Trial design</i>                                    | <i>Intervention (control = usual care unless otherwise specified)</i>  | <i>Measures with potential cost saving implications</i> | <i>Results (intervention vs. control)</i>                                    | <i>Direct measurement of cost</i> | <i>Integrated with primary clinical information system</i> | <i>Statistically &amp; clinically significant improvement in cost or proxy metric</i> | <i>Cost effectiveness study</i> |
|-------------------------|----------------------------------|-----------------------------|--|--|---|--|-----------------------------------|--|---|---------------------------------|
| Bates DW (1999) [19]    | Academic hospital                | 1 Hospital; 1817 admissions | Time series: 1 baseline period, 3 intervention periods | CPOE + CDS with serial improvements of CDS across study periods  | Non-missed-dose medication error rate                   | 26.6 (final period) vs. 142 events/1000 patient days (P < 0.0001)            | No                                | Yes  | Yes   | No                              |
|                         |                                  |                             |  |  | Non-intercepted serious medication error rate           | 1.1 (final period) vs. 7.6 events/1000 patient days (P < 0.0003)             | No                                | NR   | No  | No                              |
| East TD (1999) [20]     | Academic and community hospitals | 10 hospitals; 200 patients  | RCT  | CDS system for management of mechanical ventilation  | ICU length of stay                                      | 27.6 vs. 25.2 days (NS)  | No                                | NR   | No  | No                              |
| Durieux P (2000) [21]   | Teaching hospital                | 1 hospital; 1971 patients   | Interrupted time series                                | CDS system for provision of pharmacological VTE prophylaxis recommendations to orthopedic providers for post-operative patients                  | Rate of compliance with VTE prophylaxis guidelines      | 94.9% (95% CI: 92.5%-96.6%) vs. 82.8% (95% CI: 77.6%-87.1%)                  | No                                | Yes  | Yes   | No                              |
| Schriger DL (2000) [22] | Academic emergency department    | 1 hospital; 830 patients    | Interrupted time series: off-on-off                    | Guideline-driven electronic charting system that provided recommendations for evaluation and treatment of febrile children less than 3 years old | Median charges  | \$216 (intervention) vs. \$216 (baseline) vs. \$222 (post-intervention) (NS) | No                                | No   | No  | No                              |
|                         |                                  |                             |  |  | Mean charges  | \$387 (intervention) vs. \$357 (baseline) vs. \$635 (post-intervention) (NS) |                                   |  |   |                                 |

Table 1 Continued

| <i>Citation</i>        | <i>Setting</i>     | <i>Scope</i>               | <i>Trial design</i>     | <i>Intervention (control = usual care unless otherwise specified)</i>   | <i>Measures with potential cost saving implications</i>  | <i>Results (intervention vs. control)</i>  | <i>Direct measurement of cost</i> | <i>Integrated with primary clinical information system</i> | <i>Statistically &amp; clinically significant improvement in cost or proxy metric</i> | <i>Cost effectiveness study</i> |
|------------------------|--------------------|----------------------------|-------------------------|---|--|--|-----------------------------------|--|---|---------------------------------|
| Chertow GM (2001) [23] | Teaching hospital  | 1 hospital; 17828 patients | Interrupted time series | CPOE with CDS that provided adjusted default dose and frequency for patients with renal insufficiency                                     | Mean length of stay<br><br>Mean total costs  | 4.3 vs. 4.5 days (P = 0.009)<br><br>\$4881 vs. \$4968 (NS)   | Yes                               | Yes  | No  | No                              |
| Dexter PR (2001) [24]  | Teaching hospital  | 1 hospital; 6371 patients  | RCT                     | Computer generated reminders of patient eligibility for preventive care measures including prophylactic heparin use in high risk patients | Rate of prophylactic heparin use in high risk patients   | 32.3% vs. 18.9% (P < 0.001)  | No                                | Yes  | Yes   | No                              |
| Kellett J (2001) [25]  | Community hospital | 1 hospital; 894 patients   | Pre-post comparison     | CDS system designed to aid in the decision to treat cases of acute myocardial infarction with fibrinolytic therapy                        | Proportion of appropriate candidates who received fibrinolytic therapy   | 68.9% vs. 66.7% (NS)   | No                                | No   | No  | No                              |
| Mullett CJ (2001) [26] | Teaching hospital  | 1 hospital; 1758 patients  | Pre-post comparison     | Comprehensive antimicrobial management system in a pediatric ICU setting  | Mean hospital costs per patient<br><br>Total anti-infectives cost<br><br>Total PICU anti-infectives cost<br><br>PICU length of stay<br><br>Hospital length of stay | \$28257 vs. \$25032 (NS)<br><br>\$289.60 vs. \$274.79 (NS)<br><br>\$183.53 vs. \$177.03 (NS)<br><br>4.90 days vs. 4.93 days (NS)<br><br>10.76 vs 10.76 days (NS) | Yes                               | Yes  | No  | No                              |

Table 1 Continued

| <i>Citation</i>            | <i>Setting</i>                | <i>Scope</i>                                  | <i>Trial design</i>     | <i>Intervention (control = usual care unless otherwise specified)</i>   | <i>Measures with potential cost saving implications</i>   | <i>Results (intervention vs. control)</i>  | <i>Direct measurement of cost</i> | <i>Integrated with primary clinical information system</i> | <i>Statistically &amp; clinically significant improvement in cost or proxy metric</i> | <i>Cost effectiveness study</i> |
|----------------------------|-------------------------------|---|-------------------------|---|---|--|-----------------------------------|--|---|---------------------------------|
| Mekhjian HS (2002) [27]    | Academic hospital system      | 2 hospitals; 28898 patients                   | Pre-post comparison     | Comprehensive CPOE system with CDS for pharmacotherapy  | Severity-adjusted total cost per admission at hospital #1<br><br>Severity-adjusted total cost per admission at hospital #2                      | \$5661 vs. \$5697 (NS)   | Yes                               | Yes  | No  | No                              |
|                            |                               |   |                         |   | Severity-adjusted length of stay at hospital #1<br><br>Severity-adjusted length of stay at hospital #2  | 3.71 vs. 3.91 days (P < 0.002)   |                                   |  |   |                                 |
|                            |                               |   |                         |   | Severity-adjusted length of stay at hospital #2   | 3.61 vs. 3.68 (NS)   |                                   |  |   |                                 |
| Buller-Close K (2003) [28] | Academic emergency department | 1 hospital; 280 patients                      | Interrupted time series | Guideline-driven electronic charting system that provided recommendations for evaluation and treatment of occupational exposure to blood or body fluids         | Median charges per patient  | \$392 (intervention) vs. \$473 (baseline) (95% CI \$51 to \$161 difference in charges)   | No                                | No   | Yes   | No                              |
| Fischer MA (2003) [29]     | Academic hospital             | 1 hospital; 1045 orders in intervention phase | Pre-post comparison     | CDS within CPOE system to identify and inform providers of patients who are eligible to transition from intravenous to oral route for five targeted medications | Change in IV defined daily dose<br><br>Change in PO defined daily dose<br><br>Change in length of stay<br><br>Change in total drug expenditures | -11.1% (P = 0.002)<br><br>+3.7% (P = 0.002)<br><br>+1.9% (P = N/A)<br><br>+12% (P = N/A) | Yes                               | Yes  | No  | No                              |



Table 1 Continued

| Citation                | Setting           | Scope                                | Trial design        | Intervention (control = usual care unless otherwise specified)  | Measures with potential cost saving implications   | Results (intervention vs. control)  | Direct measurement of cost | Integrated with primary clinical information system | Statistically & clinically significant improvement in cost or proxy metric | Cost effectiveness study |
|-------------------------|-------------------|--------------------------------------|---------------------|---|--|---|----------------------------|---|--|--------------------------|
| Galanter WL (2004) [30] | Academic hospital | 1 hospital; 1596 alerting situations | Pre-post comparison | CDS designed to improve the safe use of digoxin by providing alerts related to digoxin, potassium, and magnesium serum levels | Unknown serum values checked within one hour<br><br>Unknown serum values checked within 24 hours<br><br>Low serum values supplemented within one hour<br><br>Low serum values supplemented within 24 hours | Improved with P < 0.01 for 3 of 3 measures<br><br>Improved with P < 0.01 for 3 of 3 measures<br><br>Improved with P < 0.05 for 3 of 4 measures<br><br>Improved with P < .05 for 2 of 4 measures | No                         | Yes   | Yes  | No                       |
| Hulgan T (2004) [31]    | Academic hospital | 1 hospital; 15194 orders             | Pre-post comparison | CDS to promote the use of oral—versus IV—quinolones   | Percent of quinolone orders for oral route   | 62.4% vs. 55.5% (P <= 0.001 )   | No                         | Yes   | Yes  | No                       |
| Potts AL (2004) [32]    | Academic hospital | 1 hospital; 514 patients             | Pre-post comparison | CPOE with CDS in a pediatric ICU setting  | Potential ADE rate<br><br>Medication prescribing error rate<br><br>Rule violation rate   | 1.3 vs. 2.2 events per 100 orders (P <0.001)<br><br>0.2 vs. 30.1 events per 100 orders (P < 0.001)<br><br>0.1 vs. 6.8 events per 100 orders (P < 0.001)   | No                         | Yes   | Yes  | No                       |

Table 1 Continued

| <i>Citation</i>         | <i>Setting</i>     | <i>Scope</i>                                  | <i>Trial design</i>                                    | <i>Intervention (control = usual care unless otherwise specified)</i>  | <i>Measures with potential cost saving implications</i>   | <i>Results (intervention vs. control)</i>                        | <i>Direct measurement of cost</i> | <i>Integrated with primary clinical information system</i> | <i>Statistically &amp; clinically significant improvement in cost or proxy metric</i> | <i>Cost effectiveness study</i> |
|-------------------------|--------------------|---|--|--|---|--|-----------------------------------|--|---|---------------------------------|
| Galanter WL (2005) [33] | Academic hospital  | 1 hospital; 410 alerts or alerting situations | Pre-post comparison                                    | CDS system designed to alert providers attempting to order medication contraindicated for patients with renal insufficiency  | Likelihood of patient receiving at least one dose of contraindicated medication                       | 47% vs. 89% (p < 0.0001))  | No                                | Yes  | Yes   | No                              |
| Kucher N (2005) [34]    | Academic hospital  | 1 hospital; 2506 patients                     | RCT  | CDS system to identify patients at risk for deep venous thrombosis and alert physicians  | Rate of venous thromboembolism at 90 days<br><br>Rate of pulmonary embolism at 90 days                | 4.9% vs. 8.2% (P = 0.001)<br><br>1.1% vs. 2.8% (P = 0.004)       | No                                | Yes  | Yes   | No                              |
| Nash IS (2005) [35]     | Academic hospital  | 1 hospital                                    | Time series: 1 baseline period, 2 intervention periods | CDS to identify patients with renal impairment who had received excessive dosage of medication. Physicians were then notified by nursing staff in one intervention period and notified by pharmacists in the other intervention period | Rate of excessive dosing with nurse feedback<br><br>Rate of excessive dosing with pharmacist feedback | 17.3% vs. 23.2% (p < 0.05)<br><br>16.8% vs. 23.2% (p < 0.05)     | No                                | Yes  | Yes   | No                              |
| Chan AL (2006) [36]     | Community hospital | 1 hospital                                    | Pre-post comparison                                    | CDS consisting of an automated gentamicin dosing calculator integrated into an ICU CPOE system   | Frequency of undesirable serum gentamicin levels  | 13.5% vs. 32.7% (p < 0.05)*                                      | No                                | Yes  | Yes   | No                              |
| McGregor JC (2006) [37] | Academic hospital  | 1 hospital                                    | RCT  | CDS to provide antimicrobial management support to the institution's antimicrobial management team   | Average antimicrobial expenditures per patient<br><br>Length of stay                                  | \$127.77 vs. \$163.00 (p < 0.05)*<br><br>3.84 vs. 3.99 days (NS) | Yes                               | Yes  | Yes   | No                              |

Table 1 Continued

| Citation                       | Setting            | Scope                       | Trial design        | Intervention (control = usual care unless otherwise specified)  | Measures with potential cost saving implications  | Results (intervention vs. control)        | Direct measurement of cost | Integrated with primary clinical information system | Statistically & clinically significant improvement in cost or proxy metric | Cost effectiveness study |
|--------------------------------|--------------------|-----------------------------|---------------------|---|---|---|----------------------------|---|--|--------------------------|
| Thursky KA (2006) [38]         | Community hospital | 1 hospital; 1060 admissions | Pre-post comparison | CDS for antibiotic management in an ICU setting   | Odds patient prescribed carbapenems after adjusting for risk factors                                  | OR = 0.61, 95% CI = 0.39-0.97             | No                         | Yes   | Yes  | No                       |
|                                |                    |                             |                     |   | Odds patient prescribed 3 <sup>rd</sup> generation cephalosporin after adjusting for risk factors     | OR = 0.58, 95% CI = 0.42-0.79             |                            |   |  |                          |
|                                |                    |                             |                     |   | Odds patient prescribed vancomycin after adjusting for risk factors                                   | OR = 0.67, 95% CI = 0.45-1.00             |                            |   |  |                          |
| Fernandez Perez ER (2007) [39] | Academic hospital  | 1 hospital; 2200 patients   | Pre-post comparison | CDS for management of RBC transfusions in the ICU setting   | RBC transfusion cost  | \$556226 vs. \$616442 (P = N/A)           | Yes                        | Yes   | Yes  | No                       |
|                                |                    |                             |                     |   | Average number of units transfused per patient  | 1.3 vs. 1.5 units (P = 0.045)             |                            |   |  |                          |
| Rothschild JM (2007) [40]      | Academic hospital  | 1 hospital; 1607 patients   | RCT                 | CDS for management of blood product transfusions  | Percent of transfusion orders which were not guideline compliant (typically unnecessary or excessive) | 59.6% vs. 67.5% (P < 0.0001)              | No                         | Yes   | Yes  | No                       |
| Vardi A (2007) [41]            | Academic hospital  | 1 hospital; 60094 orders    | Pre-post comparison | CDS designed to prevent medication errors for resuscitation drugs used in a pediatric critical care setting | Number of medication order errors   | 0 vs. 3 medication order errors (P = N/A) | No                         | No  | No   | No                       |

Table 1 Continued

| <i>Citation</i>         | <i>Setting</i>     | <i>Scope</i>               | <i>Trial design</i>                                    | <i>Intervention (control = usual care unless otherwise specified)</i>   | <i>Measures with potential cost saving implications</i>  | <i>Results (intervention vs. control)</i>  | <i>Direct measurement of cost</i> | <i>Integrated with primary clinical information system</i> | <i>Statistically &amp; clinically significant improvement in cost or proxy metric</i> | <i>Cost effectiveness study</i> |
|-------------------------|--------------------|----------------------------|--|---|--|--|-----------------------------------|--|---|---------------------------------|
| Buisson KL (2008) [42]  | Teaching hospital  | 1 hospital; 525 patients   | Pre-post comparison                                    | CDS designed to support adherence to community acquired pneumonia recommendations   | Percent of antibiotic prescriptions concordant with recommendations<br><br>Average cost of antibiotics per patient                           | 89.7% vs. 61.9% (P < 0.01)<br><br>\$84.04 vs. \$72.07 (P = N/A)                                    | Yes                               | Yes  | No  | No                              |
| Lecumberrir (2008) [43] | Academic hospital  | 1 hospital; 19338 patients | Time series: 1 baseline period, 2 intervention periods | CDS system for identification of patients at high risk for VTE and notification of physicians of the need for prophylaxis | VTE rate for 1 <sup>st</sup> intervention period<br><br>VTE rate for 2 <sup>nd</sup> intervention period                                     | 1.74 vs. 3.26 events/1000 patients (P < 0.05)<br><br>1.67 vs. 3.26 events/1000 patients (P < 0.05) | No                                | Yes  | Yes   | No                              |
| Roukema J (2008) [44]   | Academic hospital  | 1 hospital; 164 patients   | RCT  | CDS system for the evaluation/management of children with fever without apparent cause in the ED setting                  | Median length of ED stay<br><br>Frequency of lab orders  | 138 vs. 123 minutes (NS)<br><br>82% vs. 44% (P < 0.001)  | No                                | No   | No  | No                              |
| Eden A (2009) [45]      | Community hospital | 1 hospital; 995 patients   | Time series: 1 baseline period, 2 intervention periods | CDS system designed to remind anesthesiologists to reactivate alarms following cardiopulmonary bypass                     | Rate of alarm reactivation for 1 <sup>st</sup> intervention period<br><br>Rate of alarm reactivation for 2 <sup>nd</sup> intervention period | 63% vs. 22% (P < 0.001)<br><br>83% vs. 22% (P < 0.001)   | No                                | Yes  | Yes   | No                              |

Table 1 Continued

| Citation                | Setting            | Scope                                 | Trial design        | Intervention (control = usual care unless otherwise specified)  | Measures with potential cost saving implications  | Results (intervention vs. control)   | Direct measurement of cost | Integrated with primary clinical information system | Statistically & clinically significant improvement in cost or proxy metric | Cost effectiveness study |
|-------------------------|--------------------|---------------------------------------|---------------------|---|---|--|----------------------------|---|--|--------------------------|
| Flanders SJ (2009) [46] | Academic hospitals | 2 hospitals; 1482582 glucose measures | Pre-post comparison | CDS system designed to prevent hyperglycemia in patients receiving IV insulin in the ICU setting                                      | Odds of a glucose measure < 150 mg/dL during 1 <sup>st</sup> year post-baseline<br><br>Odds of a glucose measure < 150 mg/dL during 2 <sup>nd</sup> year post-baseline<br><br>Odds of a glucose measure < 150 mg/dL during 3 <sup>rd</sup> year post-baseline | OR = 1.8, 95% CI = 1.78-1.82<br><br>OR = 2.16, 95% CI = 2.14-2.19<br><br>OR = 2.28, 95% CI = 2.25-2.30 | No                         | No  | Yes  | No                       |
| Matsumura Y (2009) [47] | Academic hospital  | 1 hospital; 970 patients              | Pre-post comparison | CDS implemented with CPOE to prevent the ordering of contraindicated medications for patients with renal insufficiency                | Rate of discontinuation of contraindicated medication   | 54% vs. 24% (P = 0.01)   | No                         | Yes   | Yes  | No                       |
| Niemi K (2009) [48]     | Community hospital | 1 hospital; 4090 patients             | Pre-post comparison | CDS system for identification of patients with pneumonia or heart failure and recommendations to providers to meet quality indicators | Compliance with quality indicators for pneumonia and heart failure cases  | One of six measures improved with statistical significance   | No                         | Yes   | No   | No                       |

Table 1 Continued

| <i>Citation</i>        | <i>Setting</i>    | <i>Scope</i>                 | <i>Trial design</i> | <i>Intervention (control = usual care unless otherwise specified)</i>  | <i>Measures with potential cost saving implications</i>                              | <i>Results (intervention vs. control)</i> | <i>Direct measurement of cost</i> | <i>Integrated with primary clinical information system</i> | <i>Statistically &amp; clinically significant improvement in cost or proxy metric</i> | <i>Cost effectiveness study</i> |
|------------------------|-------------------|------------------------------|---------------------|--|--|---|-----------------------------------|--|---|---------------------------------|
| Riggio JM (2009) [49]  | Academic hospital | 1 hospital; 65604 admissions | Pre-post comparison | CDS system to assist in the early detection of HIT   | Average time from fall in platelet count to heparin-induced thrombocytopenia testing | 3.0 vs. 2.3 days (NS)                     | No                                | Yes  | No  | No                              |
|                        |                   |                              |                     |  | Average time from fall in platelet count to heparin-induced thrombocytopenia therapy | 15.0 vs. 19.3 days (NS)                   |                                   |  |   |                                 |
|                        |                   |                              |                     |  | Average time from fall in platelet count to discontinuation of heparin products      | 2.9 vs. 1.3 days ( $p = 0.04$ )           |                                   |  |   |                                 |
| Sellier E (2009) [50]  | Teaching hospital | 1 hospital; 603 patients     | Pre-post comparison | CPOE with CDS intended to decrease the proportion of inappropriate prescriptions in patients with renal insufficiency  | Proportion of inappropriate prescriptions  | 19.9% vs. 21.3% (NS)                      | No                                | Yes  | No  | No                              |
| Bertsche T (2010) [51] | Academic hospital | 1 hospital; 265 patients     | Pre-post comparison | CDS system designed to decrease the incidence of ADEs related to drug-drug interactions in patients prescribed >8 drugs in the ICU setting. Printed alerts were provided to physicians | Relative risk of patient experiencing at least one ADE due to drug-drug interactions | 25% vs. 44% ( $P < 0.01$ )                | No                                | No   | Yes   | No                              |

Table 1 Continued

| <i>Citation</i>         | <i>Setting</i>    | <i>Scope</i>               | <i>Trial design</i> | <i>Intervention (control = usual care unless otherwise specified)</i>  | <i>Measures with potential cost saving implications</i>   | <i>Results (intervention vs. control)</i>                              | <i>Direct measurement of cost</i> | <i>Integrated with primary clinical information system</i> | <i>Statistically &amp; clinically significant improvement in cost or proxy metric</i> | <i>Cost effectiveness study</i> |
|-------------------------|-------------------|----------------------------|---------------------|--|---|--|-----------------------------------|--|---|---------------------------------|
| Elkin PL (2010) [52]    | Academic hospital | 1 hospital; 1737 cases     | Pre-post comparison | Computer-based diagnostic decision support system made available to residents in a teaching hospital   | Mean cost per diagnostically challenging case<br><br>Mean total charges per diagnostically challenging case | \$7382 vs. \$8318 (P = 0.0054)<br><br>\$11403 vs. \$12684 (P = 0.0180) | Yes                               | No   | Yes   | Yes                             |
|                         |                   |                            |                     |  | Mean Medicare Part A charges per diagnostically challenging case  | \$9390 vs. \$10422 (P = 0.0173)  |                                   |  |   |                                 |
|                         |                   |                            |                     |  | Length of Stay  | 3.99 vs. 4.14 days (NS)  |                                   |  |   |                                 |
| Galanter WL (2010) [53] | Academic hospital | 1 hospital; 38647 patients | Pre-post comparison | Mandatory VTE risk assessment form integrated into the hospital's CPOE system. System provided recommendations for pharmacological prophylaxis based on risk | Overall VTE rates   | 0.43% vs 0.51% (NS)  | No                                | Yes  | No  | No                              |
| McCoy AB (2010) [54]    | Academic hospital | 1 hospital; 1659 patients  | Pre-post comparison | CDS system integrated in CPOE system to improve medication ordering for patients with acute kidney injury  | Rate of modification or discontinuation of contraindicated drugs  | 52.6 vs. 35.2 /100 events (P < 0.001)                                  | No                                | Yes  | Yes   | No                              |

Table 1 Continued

| Citation               | Setting           | Scope                       | Trial design        | Intervention (control = usual care unless otherwise specified)   | Measures with potential cost saving implications   | Results (intervention vs. control)   | Direct measurement of cost | Integrated with primary clinical information system | Statistically & clinically significant improvement in cost or proxy metric | Cost effectiveness study |
|------------------------|-------------------|-----------------------------|---------------------|--|--|--|----------------------------|---|--|--------------------------|
| Roberts GW (2010) [55] | Teaching hospital | 1 hospital; 1001 patients   | Pre-post comparison | CDS system for recommendations on appropriate prescription of certain renally cleared drugs in patients with renal insufficiency | Enoxaparin dosing concordant with guideline<br><br>Gentamicin dosing concordant with guideline<br><br>Vancomycin dosing concordant with guideline<br><br>Rate of holding renally cleared drugs during periods of acute renal failure | 86% vs. 68% (P = 0.03)<br><br>87% vs. 63% (P = 0.01)<br><br>77% vs. 47% (NS)<br><br>62% vs. 38% (p = 0.01) | No                         | Yes   | Yes  | No                       |
| Tafelski S (2010) [56] | Academic hospital | 1 hospital; 186 patients    | Pre-post comparison | CDS intended to assist in the management of sepsis and septic shock in an ICU setting  | Concordance with sepsis management guidelines<br><br>Mean daily antibiotic usage<br><br>Percent of ICU days which were antibiotic-free   | 90.8% vs. 39.8% (P < 0.05)<br><br>1.3 vs. 1.5 agents/day (P < 0.05)<br><br>25% vs. 18.4% (P < 0.05)        | No                         | No  | Yes  | No                       |
| Adams ES (2011) [57]   | Academic hospital | 1 hospital; 6785 discharges | Pre-post comparison | CDS system intended to support the management of children potentially needing RBC transfusions                                   | Rate of RBC transfusions per patient<br><br>Estimated direct cost savings during intervention phase  | 0.05 vs. 0.076 (P < 0.03)<br><br>\$165000 (P = N/A)  | No                         | Yes   | Yes  | No                       |



Table 1 Continued

| <i>Citation</i>         | <i>Setting</i>    | <i>Scope</i>              | <i>Trial design</i> | <i>Intervention (control = usual care unless otherwise specified)</i> | <i>Measures with potential cost saving implications</i>  | <i>Results (intervention vs. control)</i> | <i>Direct measurement of cost</i> | <i>Integrated with primary clinical information system</i> | <i>Statistically &amp; clinically significant improvement in cost or proxy metric</i> | <i>Cost effectiveness study</i> |
|-------------------------|-------------------|---------------------------|---------------------|---|--|---|-----------------------------------|--|---|---------------------------------|
| Austrian JS (2011) [58] | Academic hospital | 1 hospital; 2087 patients | Pre-post comparison | CDS system to assist in the early detection of HIT                    | Proportion of patients tested for HIT antibody within 24 hours of a qualifying drop in platelets           | 41.9% vs. 31.5% (p = 0.03)                | No                                | Yes  | Yes   | No                              |
|                         |                   |                           |                     |   | Proportion of patients for whom heparin was discontinued within 24 hours of a qualifying drop in platelets | 26.5% vs. 21.2% (p = 0.01)                |                                   |  |   |                                 |
|                         |                   |                           |                     |   | Proportion of patients above the median length of stay adjusted for comorbidities                          | 50.3 vs. 49.7 (NS)                        |                                   |  |   |                                 |
| Cox ZL (2011) [59]      | Academic hospital | 1 hospital; 216 patients  | Pre-post comparison | CDS system to assist in the prescription of tobramycin and amikacin   | Proportion of initial doses within 10% of the calculated recommended dose                                  | 80% vs. 40% (P < 0.001)                   | No                                | Yes  | Yes   | No                              |
|                         |                   |                           |                     |   | Percentage of correct initial dosing interval  | 87% vs. 63% (P < 0.001)                   |                                   |  |   |                                 |
|                         |                   |                           |                     |   | Incidence of nephrotoxicity  | 17% vs. 25% (NS)                          |                                   |  |   |                                 |

Table 1 Continued

| <i>Citation</i>          | <i>Setting</i>      | <i>Scope</i>              | <i>Trial design</i>        | <i>Intervention (control = usual care unless otherwise specified)</i>  | <i>Measures with potential cost saving implications</i>                                 | <i>Results (intervention vs. control)</i>  | <i>Direct measurement of cost</i> | <i>Integrated with primary clinical information system</i> | <i>Statistically &amp; clinically significant improvement in cost or proxy metric</i> | <i>Cost effectiveness study</i> |
|--------------------------|---------------------|---------------------------|----------------------------|--|---|--|-----------------------------------|--|---|---------------------------------|
| Fitzgerald M (2011) [60] | Teaching hospital   | 1 hospital; 1171 cases    | RCT                        | CDS system to promote adherence to trauma protocols in the ED setting  | Hospital length of stay<br><br>Average ICU length of stay<br><br>Error rate per patient | NS<br><br>70 vs. 112 hours (NS)<br><br>2.13 vs. 2.30 errors per patient (P = 0.04) | No                                | No   | Yes   | No                              |
|                          |                     |                           |                            |  | Percent of patients who received packed red blood cells                                 | 66.1% vs. 84.9% (P < 0.001)  |                                   |  |   |                                 |
| Giuliano KK (2011) [61]  | Community hospitals | 2 hospitals; 135 patients | Pre-post comparison        | CDS system designed to increase adherence to sepsis protocols in the ICU setting   | Total hospital length of stay<br><br>ICU length of stay                                 | 17.8 vs. 15.7 days (NS)<br><br>10.3 vs. 7.4 days (NS)                              | No                                | No   | No  | No                              |
| Kazemi A (2011) [62]     | Teaching hospital   | 1 hospital; 248 patients  | Pre-post comparison        | CDS implemented within a CPOE system in a neonatal ward setting. CDS was specifically designed to address medication dosing errors | Medication error rate   | 34% vs. 53% (P < 0.001)  | No                                | Yes  | Yes   | No                              |
| Lipton JA (2011) [63]    | Academic hospital   | 1 hospital; 667 patients  | Pre-post comparison        | CDS system intended to assist with the management of intravenous insulin therapy in a coronary critical care unit                  | Proportion of patients with a mean blood glucose level within the target range          | 43% vs. 31% (P = 0.01)   | No                                | Yes  | Yes   | No                              |
| Mann EA (2011) [64]      | Teaching hospital   | 1 hospital; 18 patients   | Randomized crossover trial | CDS intended to support the management of insulin in burn patients in the ICU setting  | Percent of time in target glucose range   | 47% vs. 41% (p < 0.05)   | No                                | No   | Yes   | No                              |

Table 1 Continued

| <i>Citation</i>       | <i>Setting</i>    | <i>Scope</i>             | <i>Trial design</i>            | <i>Intervention (control = usual care unless otherwise specified)</i>  | <i>Measures with potential cost saving implications</i>  | <i>Results (intervention vs. control)</i>   | <i>Direct measurement of cost</i> | <i>Integrated with primary clinical information system</i> | <i>Statistically &amp; clinically significant improvement in cost or proxy metric</i> | <i>Cost effectiveness study</i> |
|-----------------------|-------------------|--------------------------|--------------------------------|--|--|---|-----------------------------------|--|---|---------------------------------|
| Milani RV (2011) [65] | Academic hospital | 1 hospital; 80 patients  | Prospectively controlled trial | CPOE with CDS to assist the prescription of antithrombotics in patients with chronic kidney disease admitted for acute coronary syndrome         | Proportion of patients who received contraindicated antithrombotics<br><br>Proportion of patients who experienced in-hospital bleeding | 0% vs 17% (P = 0.01)<br><br>9% vs. 21% (NS) | No                                | Yes  | Yes   | No                              |
|                       |                   |                          |                                |  | Length of stay   | 4.8 vs. 9.1 days (P = 0.01)                 |                                   |  |   |                                 |
| Qian Q (2011) [66]    | Academic hospital | 1 hospital               | Pre-post comparison            | CDS system intended to alert physicians to patients with left ventricular systolic dysfunction who were not being treated with an ACEi/ARB agent | Percentage of ACEi/ARB adherence   | 97.6% vs. 88.4% (P < 0.01)                  | No                                | Yes  | Yes   | No                              |
| Salinas J (2011) [67] | Teaching hospital | 1 hospital; 105 patients | Pre-post comparison            | CDS system designed to support the fluid resuscitation of severely burned patients in the ICU setting  | Mean ventilator free days<br><br>Mean ICU free days  | 6.5 vs. 3.8 days (P < 0.05)<br><br>NS       | No                                | No   | Yes   | No                              |

Table 1 Continued

| <i>Citation</i>        | <i>Setting</i>    | <i>Scope</i>             | <i>Trial design</i>     | <i>Intervention (control = usual care unless otherwise specified)</i>  | <i>Measures with potential cost saving implications</i>   | <i>Results (intervention vs. control)</i>   | <i>Direct measurement of cost</i> | <i>Integrated with primary clinical information system</i> | <i>Statistically &amp; clinically significant improvement in cost or proxy metric</i> | <i>Cost effectiveness study</i> |
|------------------------|-------------------|--------------------------|-------------------------|--|---|---|-----------------------------------|--|---|---------------------------------|
| Cho A (2012) [68]      | Academic hospital | 1 hospital; 463 patients | Pre-post comparison     | CDS system intended to alert physicians to patients at high risk for contrast-induced AKI when ordering contrast-enhanced CT imaging | Percent of at-risk patients who received prophylaxis for contrast-induced AKI<br><br>Incidence of contrast-induced AKI among at-risk patients | 55% vs. 25% (P < 0.001)<br><br>3% vs. 10% (P < 0.02)                                | No                                | Yes  | Yes   | No                              |
| Griffey RT (2012) [69] | Academic ED       | 1 ED; 1407 patients      | Interrupted time-series | CDS system designed to guide medication dosing in geriatric patients in an ED setting  | Proportion of orders consistent with dosing recommendations<br><br>Proportion of patients who experienced an ADE                              | 31.4% vs. 23% (P < 0.001)<br><br>3.4% vs. 7.1% (P = 0.02)                           | No                                | Yes  | Yes   | No                              |
| Raja AS (2012) [70]    | Academic ED       | 1 ED; 6838 patients      | Pre-post comparison     | CDS system intended to assist with the use of CT pulmonary angiography for acute pulmonary embolism                                  | ED length of stay<br><br>Quarterly utilization of CT pulmonary angiography  | 5.6 vs. 5.8 hours (NS)<br><br>21.1 vs. 26.4 examinations/1000 patients (P = 0.0379) | No                                | No   | Yes   | No                              |

Table 1 Continued

| Citation           | Setting           | Scope                           | Trial design        | Intervention (control = usual care unless otherwise specified)  | Measures with potential cost saving implications  | Results (intervention vs. control) | Direct measurement of cost | Integrated with primary clinical information system | Statistically & clinically significant improvement in cost or proxy metric | Cost effectiveness study |
|--------------------|-------------------|---------------------------------|---------------------|---|---|------------------------------------|----------------------------|---|--|--------------------------|
| Wang H (2012) [71] | Teaching hospital | 1 hospital; 38647 prescriptions | Pre-post comparison | CDS within CPOE system intended to prevent dosing errors of 13 antibiotics which require adjustment based on creatinine clearance | Percent of antibiotic prescriptions with an inappropriate dosage during the 4 <sup>th</sup> year following implementation | 3.42% vs. 21.3% (p < 0.001)        | No                         | Yes   | Yes  | No                       |
|                    |                   |                                 |                     |   | Incidence rate of renal function deterioration during the 4 <sup>th</sup> year following implementation                   | 9.47% vs. 10.94 (p < 0.001)        |                            |   |  |                          |

ACEi = angiotensin-converting-enzyme inhibitor; ADE = adverse drug event; AKI = acute kidney injury; ARB = angiotensin receptor blocker; CDS = clinical decision support; CPOE = computerized provider order entry; CT = computed tomography; ED = emergency department; HIT = heparin-induced thrombocytopenia; ICU = intensive care unit; NR = not reported; NS = not significant; OR = odds ratio; RBC = red blood cell; RCT = randomized controlled trial; VTE = venous thromboembolism

\*p value not provided by manuscript; calculated using raw data provided in manuscript

#p value not provided by manuscript; calculated using raw data provided in manuscript; assumes historical control period had same sample size as intervention period

## **CONCLUSIONS**

Health IT, and CDS in particular, has been touted for many years as a highly promising strategy for improving clinical care and “bending the cost curve” [78, 79]. However, more recent analyses have found that health IT systems such as EHR systems are not having the anticipated benefits in cost reduction [80, 81]. This study adds to these concerns that the potential benefits of health IT and CDS are not well grounded in empirical evidence, with only nine studies directly measuring costs and only one actually measuring cost-effectiveness of CDS for inpatient cost reduction.

As healthcare organizations continue to rapidly adopt health IT, leadership within those organizations must decide how to best use limited resources. Presumably, the potential cost savings associated with intervention candidates is a major factor in making those decisions. However, as a discipline, informatics does not appear to be meeting the needs of these healthcare decision makers with regard to CDS, as we have not been providing sufficient, rigorous data related to the cost benefits of CDS interventions in the inpatient setting. Further research with specific attention to cost implications of CDS systems is clearly needed.

## **APPENDIX**

Detailed search strategy:

1. exp Decision Support Systems, Clinical/
2. Decision Making, Computer-Assisted/
3. exp Reminder Systems/
4. exp Guideline Adherence/
5. exp Medical Informatics/
6. 4 and 5
7. 1 or 2 or 3 or 6
8. limit 7 to (English language and humans)

## REFERENCES

1. Keehan SP, Cuckler GA, Sisko AM, Madison AJ, Smith SD, Lizonitz JM, Poisal JA, Wolfe CJ: **National health expenditure projections: modest annual growth until coverage expands and economic growth accelerates.** *Health Aff (Millwood)* 2012, **31**(7):1600-1612.
2. Miller HD: **From volume to value: better ways to pay for health care.** *Health Aff (Millwood)* 2009, **28**(5):1418-1428.
3. Osheroff JA, Teich JM, Middleton B, Steen EB, Wright A, Detmer DE: **A roadmap for national action on clinical decision support.** *J Am Med Inform Assoc* 2007, **14**(2):141-145.
4. Hunt DL, Haynes RB, Hanna SE, Smith K: **Effects of computer-based clinical decision support systems on physician performance and patient outcomes: a systematic review.** *JAMA* 1998, **280**(15):1339-1346.
5. Kaushal R, Shojania KG, Bates DW: **Effects of computerized physician order entry and clinical decision support systems on medication safety: a systematic review.** *Arch Intern Med* 2003, **163**(12):1409-1416.
6. Garg AX, Adhikari NK, McDonald H, Rosas-Arellano MP, Devereaux PJ, Beyene J, Sam J, Haynes RB: **Effects of computerized clinical decision support systems on practitioner performance and patient outcomes: a systematic review.** *JAMA* 2005, **293**(10):1223-1238.
7. Kawamoto K, Houlihan CA, Balas EA, Lobach DF: **Improving clinical practice using clinical decision support systems: a systematic review of trials to identify features critical to success.** *BMJ* 2005, **330**(7494):765.
8. Sintchenko V, Magrabi F, Tipper S: **Are we measuring the right end-points? Variables that affect the impact of computerised decision support on patient outcomes: a systematic review.** *Med Inform Internet Med* 2007, **32**(3):225-240.
9. Shojania KG, Jennings A, Mayhew A, Ramsay C, Eccles M, Grimshaw J: **Effect of point-of-care computer reminders on physician behaviour: a systematic review.** *CMAJ* 2010, **182**(5):E216-225.
10. Chaudhry B, Wang J, Wu S, Maglione M, Mojica W, Roth E, Morton SC, Shekelle PG: **Systematic review: impact of health information technology on quality, efficiency, and costs of medical care.** *Ann Intern Med* 2006, **144**(10):742-752.
11. Bright TJ, Wong A, Dhurjati R, Bristow E, Bastian L, Coeytaux RR, Samsa G, Hasselblad V, Williams JW, Musty MD *et al*: **Effect of clinical decision-support systems: a systematic review.** *Ann Intern Med* 2012, **157**(1):29-43.
12. Larsen RA, Evans RS, Burke JP, Pestotnik SL, Gardner RM, Classen DC: **Improved perioperative antibiotic use and reduced surgical wound infections through use of computer decision analysis.** *Infect Control Hosp Epidemiol* 1989, **10**(7):316-320.



13. Rind DM, Safran C, Phillips RS, Slack WV, Calkins DR, Delbanco TL, Bleich HL: **The effect of computer-based reminders on the management of hospitalized patients with worsening renal function.** *Proc Annu Symp Comput Appl Med Care* 1991:28-32.
14. Day F, Hoang LP, Ouk S, Nagda S, Schrager DL: **The impact of a guideline-driven computer charting system on the emergency care of patients with acute low back pain.** *Proc Annu Symp Comput Appl Med Care* 1995:576-580.
15. Bates DW, Leape LL, Cullen DJ, Laird N, Petersen LA, Teich JM, Burdick E, Hickey M, Kleeffeld S, Shea B *et al*: **Effect of computerized physician order entry and a team intervention on prevention of serious medication errors.** *JAMA* 1998, **280**(15):1311-1316.
16. Evans RS, Pestotnik SL, Classen DC, Clemmer TP, Weaver LK, Orme JF, Jr., Lloyd JF, Burke JP: **A computer-assisted management program for antibiotics and other antiinfective agents.** *N Engl J Med* 1998, **338**(4):232-238.
17. Knirsch CA, Jain NL, Pablos-Mendez A, Friedman C, Hripcsak G: **Respiratory isolation of tuberculosis patients using clinical guidelines and an automated clinical decision support system.** *Infect Control Hosp Epidemiol* 1998, **19**(2):94-100.
18. Aase O: **Clinical experience with a decision support computer program using Bayes' theorem to diagnose chest pain patients.** *Cardiology* 1999, **92**(2):128-134.
19. Bates DW, Teich JM, Lee J, Seger D, Kuperman GJ, Ma'Luf N, Boyle D, Leape L: **The impact of computerized physician order entry on medication error prevention.** *J Am Med Inform Assoc* 1999, **6**(4):313-321.
20. East TD, Heermann LK, Bradshaw RL, Lugo A, Sailors RM, Ershler L, Wallace CJ, Morris AH, McKinley B, Marquez A *et al*: **Efficacy of computerized decision support for mechanical ventilation: results of a prospective multi-center randomized trial.** *Proc AMIA Symp* 1999, **Annual Symposium**.:251-255.
21. Durieux P, Nizard R, Ravaud P, Mounier N, Lepage E: **A clinical decision support system for prevention of venous thromboembolism: effect on physician behavior.** *JAMA* 2000, **283**(21):2816-2821.
22. Schrager DL, Baraff LJ, Buller K, Shendrikar MA, Nagda S, Lin EJ, Mikulich VJ, Cretin S: **Implementation of clinical guidelines via a computer charting system: effect on the care of febrile children less than three years of age.** *J Am Med Inform Assoc* 2000, **7**(2):186-195.
23. Chertow GM, Lee J, Kuperman GJ, Burdick E, Horsky J, Seger DL, Lee R, Mekala A, Song J, Komaroff AL *et al*: **Guided medication dosing for inpatients with renal insufficiency.** *JAMA* 2001, **286**(22):2839-2844.
24. Dexter PR, Perkins S, Overhage JM, Maharry K, Kohler RB, McDonald CJ: **A computerized reminder system to increase the use of preventive care for hospitalized patients.** *N Engl J Med* 2001, **345**(13):965-970.
25. Kellett J: **Decision support and the appropriate use of fibrinolysis in myocardial infarction.** *Eff Clin Pract* 2001, **4**(1):1-9.
26. Mullett CJ, Evans RS, Christenson JC, Dean JM: **Development and impact of a computerized pediatric antiinfective decision support program.** *Pediatrics* 2001, **108**(4):E75.

27. Mekhjian HS, Kumar RR, Kuehn L, Bentley TD, Teater P, Thomas A, Payne B, Ahmad A: **Immediate benefits realized following implementation of physician order entry at an academic medical center.** *J Am Med Inform Assoc* 2002, **9**(5):529-539.
28. Buller-Close K, Schriger DL, Baraff LJ: **Heterogeneous effect of an Emergency Department Expert Charting System.** *Ann Emerg Med* 2003, **41**(5):644-652.
29. Fischer MA, Solomon DH, Teich JM, Avorn J: **Conversion from intravenous to oral medications: assessment of a computerized intervention for hospitalized patients.** *Arch Intern Med* 2003, **163**(21):2585-2589.
30. Galanter WL, Polikaitis A, DiDomenico RJ: **A trial of automated safety alerts for inpatient digoxin use with computerized physician order entry.** *J Am Med Inform Assoc* 2004, **11**(4):270-277.
31. Hulgian T, Rosenbloom ST, Hargrove F, Talbert DA, Arbogast PG, Bansal P, Miller RA, Kernodle DS: **Oral quinolones in hospitalized patients: an evaluation of a computerized decision support intervention.** *J Intern Med* 2004, **256**(4):349-357.
32. Potts AL, Barr FE, Gregory DF, Wright L, Patel NR: **Computerized physician order entry and medication errors in a pediatric critical care unit.** *Pediatrics* 2004, **113**(1 Pt 1):59-63.
33. Galanter WL, DiDomenico RJ, Polikaitis A: **A trial of automated decision support alerts for contraindicated medications using computerized physician order entry.** *J Am Med Inform Assoc* 2005, **12**(3):269-274.
34. Kucher N, Koo S, Quiroz R, Cooper JM, Paterno MD, Soukonnikov B, Goldhaber SZ: **Electronic alerts to prevent venous thromboembolism among hospitalized patients.** *N Engl J Med* 2005, **352**(10):969-977.
35. Nash IS, Rojas M, Hebert P, Marrone SR, Colgan C, Fisher LA, Caliendo G, Chassin MR: **Reducing excessive medication administration in hospitalized adults with renal dysfunction.** *Am J Med Qual* 2005, **20**(2):64-69.
36. Chan ALF, Wang H-Y, Leung HWC: **Incorporation of a gentamicin dosage calculator into a computerized prescriber-order-entry system.** *Am J Health-Syst Pharm* 2006, **63**(14):1344-1345.
37. McGregor JC, Weekes E, Forrest GN, Standiford HC, Perencevich EN, Furuno JP, Harris AD: **Impact of a computerized clinical decision support system on reducing inappropriate antimicrobial use: a randomized controlled trial.** *J Am Med Inform Assoc* 2006, **13**(4):378-384.
38. Thursky KA, Buising KL, Bak N, Macgregor L, Street AC, Macintyre CR, Presneill JJ, Cade JF, Brown GV: **Reduction of broad-spectrum antibiotic use with computerized decision support in an intensive care unit.** *Int J Qual Health Care* 2006, **18**(3):224-231.
39. Fernandez Perez ER, Winters JL, Gajic O: **The addition of decision support into computerized physician order entry reduces red blood cell transfusion resource utilization in the intensive care unit.** *Am J Hematol* 2007, **82**(7):631-633.
40. Rothschild JM, McGurk S, Honour M, Lu L, McClendon AA, Srivastava P, Churchill WH, Kaufman RM, Avorn J, Cook EF *et al*: **Assessment of education and computerized decision support interventions for improving transfusion practice.** *Transfusion* 2007, **47**(2):228-239.
41. Vardi A, Efrati O, Levin I, Matok I, Rubinstein M, Paret G, Barzilay Z: **Prevention of potential errors in resuscitation medications orders by means of a computerised physician order entry in paediatric critical care.** *Resuscitation* 2007, **73**(3):400-406.

42. Buising KL, Thursky KA, Black JF, MacGregor L, Street AC, Kennedy MP, Brown GV: **Improving antibiotic prescribing for adults with community acquired pneumonia: Does a computerised decision support system achieve more than academic detailing alone?--A time series analysis.** *BMC Med Inf Decis Mak* 2008, **8**:35.
43. Lecumberri R, Marques M, Diaz-Navarraz MT, Panizo E, Toledo J, Garcia-Mouriz A, Paramo JA: **Maintained effectiveness of an electronic alert system to prevent venous thromboembolism among hospitalized patients.** *Thromb Haemost* 2008, **100**(4):699-704.
44. Roukema J, Steyerberg EW, van der Lei J, Moll HA: **Randomized trial of a clinical decision support system: impact on the management of children with fever without apparent source.** *J Am Med Inform Assoc* 2008, **15**(1):107-113.
45. Eden A, Pizov R, Toderis L, Kantor G, Perel A: **The impact of an electronic reminder on the use of alarms after separation from cardiopulmonary bypass.** *Anesth Analg* 2009, **108**(4):1203-1208.
46. Flanders SJ, Juneja R, Roudebush CP, Carroll J, Golas A, Elias BL: **Glycemic control and insulin safety: the impact of computerized intravenous insulin dosing.** *Am J Med Qual* 2009, **24**(6):489-497.
47. Matsumura Y, Yamaguchi T, Hasegawa H, Yoshihara K, Zhang Q, Mineno T, Takeda H: **Alert system for inappropriate prescriptions relating to patients' clinical condition.** *Methods Inf Med* 2009, **48**(6):566-573.
48. Niemi K, Geary S, Quinn B, Larrabee M, Brown K: **Implementation and evaluation of electronic clinical decision support for compliance with pneumonia and heart failure quality indicators.** *Am J Health-Syst Pharm* 2009, **66**(4):389-397.
49. Riggio JM, Cooper MK, Leiby BE, Walenga JM, Merli GJ, Gottlieb JE: **Effectiveness of a clinical decision support system to identify heparin induced thrombocytopenia.** *J Thromb Thrombolysis* 2009, **28**(2):124-131.
50. Sellier E, Colombet I, Sabatier B, Breton G, Nies J, Zapletal E, Arlet J-B, Somme D, Durieux P: **Effect of alerts for drug dosage adjustment in inpatients with renal insufficiency.** *J Am Med Inform Assoc* 2009, **16**(2):203-210.
51. Bertsche T, Pfaff J, Schiller P, Kaltschmidt J, Pruszydlo MG, Stremmel W, Walter-Sack I, Haefeli WE, Encke J: **Prevention of adverse drug reactions in intensive care patients by personal intervention based on an electronic clinical decision support system.** *Intensive Care Med* 2010, **36**(4):665-672.
52. Elkin PL, Liebow M, Bauer BA, Chaliki S, Wahner-Roedler D, Bundrick J, Lee M, Brown SH, Froehling D, Bailey K *et al*: **The introduction of a diagnostic decision support system (DXplainTM) into the workflow of a teaching hospital service can decrease the cost of service for diagnostically challenging Diagnostic Related Groups (DRGs).** *Int J Med Inf* 2010, **79**(11):772-777.
53. Galanter WL, Thambi M, Rosencranz H, Shah B, Falck S, Lin F-J, Nutescu E, Lambert B: **Effects of clinical decision support on venous thromboembolism risk assessment, prophylaxis, and prevention at a university teaching hospital.** *Am J Health-Syst Pharm* 2010, **67**(15):1265-1273.
54. McCoy AB, Waitman LR, Gadd CS, Danciu I, Smith JP, Lewis JB, Schildcrout JS, Peterson JF: **A computerized provider order entry intervention for medication safety during acute kidney injury: a quality improvement report.** *Am J Kidney Dis* 2010, **56**(5):832-841.

55. Roberts GW, Farmer CJ, Cheney PC, Govis SM, Belcher TW, Walsh SA, Adams RJ: **Clinical decision support implemented with academic detailing improves prescribing of key renally cleared drugs in the hospital setting.** *J Am Med Inform Assoc* 2010, **17**(3):308-312.
56. Tafelski S, Nachtigall I, Deja M, Tamarkin A, Trefzer T, Halle E, Wernecke KD, Spies C: **Computer-assisted decision support for changing practice in severe sepsis and septic shock.** *J Int Med Res* 2010, **38**(5):1605-1616.
57. Adams ES, Longhurst CA, Pageler N, Widen E, Franzon D, Cornfield DN: **Computerized physician order entry with decision support decreases blood transfusions in children.** *Pediatrics* 2011, **127**(5):e1112-1119.
58. Austrian JS, Adelman JS, Reissman SH, Cohen HW, Billett HH: **The impact of the heparin-induced thrombocytopenia (HIT) computerized alert on provider behaviors and patient outcomes.** *J Am Med Inform Assoc* 2011, **18**(6):783-788.
59. Cox ZL, Nelsen CL, Waitman LR, McCoy JA, Peterson JF: **Effects of clinical decision support on initial dosing and monitoring of tobramycin and amikacin.** *Am J Health-Syst Pharm* 2011, **68**(7):624-632.
60. Fitzgerald M, Cameron P, Mackenzie C, Farrow N, Scicluna P, Gocentas R, Bystrzycki A, Lee G, O'Reilly G, Andrianopoulos N *et al*: **Trauma resuscitation errors and computer-assisted decision support.** *Arch Surg* 2011, **146**(2):218-225.
61. Giuliano KK, Lecardo M, Staul L: **Impact of protocol watch on compliance with the surviving sepsis campaign.** *Am J Crit Care* 2011, **20**(4):313-321.
62. Kazemi A, Ellenius J, Pourasghar F, Tofighi S, Salehi A, Amanati A, Fors UGH: **The effect of computerized physician order entry and decision support system on medication errors in the neonatal ward: experiences from an Iranian teaching hospital.** *J Med Syst* 2011, **35**(1):25-37.
63. Lipton JA, Barendse RJ, Schinkel AFL, Akkerhuis KM, Simoons ML, Sijbrands EJG: **Impact of an alerting clinical decision support system for glucose control on protocol compliance and glycemic control in the intensive cardiac care unit.** *Diabetes Technol Ther* 2011, **13**(3):343-349.
64. Mann EA, Jones JA, Wolf SE, Wade CE: **Computer decision support software safely improves glycemic control in the burn intensive care unit: a randomized controlled clinical study.** *J Burn Care Res* 2011, **32**(2):246-255.
65. Milani RV, Oleck SA, Lavie CJ: **Medication errors in patients with severe chronic kidney disease and acute coronary syndrome: the impact of computer-assisted decision support.** *Mayo Clin Proc* 2011, **86**(12):1161-1164.
66. Qian Q, Manning DM, Ou N, Klarich MJ, Leutink DJ, Loth AR, Lopez-Jimenez F: **ACEi/ARB for systolic heart failure: closing the quality gap with a sustainable intervention at an academic medical center.** *J Hosp Med* 2011, **6**(3):156-160.
67. Salinas J, Chung KK, Mann EA, Cancio LC, Kramer GC, Serio-Melvin ML, Renz EM, Wade CE, Wolf SE: **Computerized decision support system improves fluid resuscitation following severe burns: an original study.** *Crit Care Med* 2011, **39**(9):2031-2038.
68. Cho A, Lee JE, Yoon JY, Jang HR, Huh W, Kim Y-G, Kim DJ, Oh HY: **Effect of an electronic alert on risk of contrast-induced acute kidney injury in hospitalized patients undergoing computed tomography.** *Am J Kidney Dis* 2012, **60**(1):74-81.

69. Griffey RT, Lo HG, Burdick E, Keohane C, Bates DW: **Guided medication dosing for elderly emergency patients using real-time, computerized decision support.** *J Am Med Inform Assoc* 2012, **19**(1):86-93.
70. Raja AS, Ip IK, Prevedello LM, Sodickson AD, Farkas C, Zane RD, Hanson R, Goldhaber SZ, Gill RR, Khorasani R: **Effect of computerized clinical decision support on the use and yield of CT pulmonary angiography in the emergency department.** *Radiology* 2012, **262**(2):468-474.
71. Wang H-Y, Lu C-L, Wu M-P, Huang M-H, Huang Y-B: **Effectiveness of an integrated CPOE decision-supporting system with clinical pharmacist monitoring practice in preventing antibiotic dosing errors.** *Int J Clin Pharmacol Ther* 2012, **50**(6):375-382.
72. Harris AD, McGregor JC, Perencevich EN, Furuno JP, Zhu J, Peterson DE, Finkelstein J: **The use and interpretation of quasi-experimental studies in medical informatics.** *J Am Med Inform Assoc* 2006, **13**(1):16-23.
73. Liu JLY, Wyatt JC: **The case for randomized controlled trials to assess the impact of clinical information systems.** *J Am Med Inform Assoc* 2011, **18**(2):173-180.
74. Higgins JPT, Green S (editors): **Cochrane handbook for systematic reviews of interventions, version 5.1.0 [updated March 2011].** The Cochrane Collaboration, 2011. Available from <http://www.cochrane-handbook.org>.
75. Deeks JJ, Dinnes J, D'Amico R, Sowden AJ, Sakarovich C, Song F, Petticrew M, Altman DG: **Evaluating non-randomised intervention studies.** *Health technology assessment (Winchester, England)* 2003, **7**(27):iii-x, 1-173.
76. Thomas BH, Ciliska D, Dobbins M, Micucci S: **A process for systematically reviewing the literature: providing the research evidence for public health nursing interventions.** *Worldviews on evidence-based nursing / Sigma Theta Tau International, Honor Society of Nursing* 2004, **1**(3):176-184.
77. Finkler SA: **The distinction between cost and charges.** *Ann Intern Med* 1982, **96**(1):102-109.
78. Hillestad R, Bigelow J, Bower A, Girosi F, Meili R, Scoville R, Taylor R: **Can electronic medical record systems transform health care? Potential health benefits, savings, and costs.** *Health Aff (Millwood)* 2005, **24**(5):1103-1117.
79. Mongan JJ, Ferris TG, Lee TH: **Options for slowing the growth of health care costs.** *N Engl J Med* 2008, **358**(14):1509-1514.
80. Kellermann AL, Jones SS: **What it will take to achieve the as-yet-unfulfilled promises of health information technology.** *Health Aff (Millwood)* 2013, **32**(1):63-68.
81. Himmelstein DU, Wright A, Woolhandler S: **Hospital computing and the costs and quality of care: a national study.** *Am J Med* 2010, **123**(1):40-46.